



Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com



Recommendations

Respiratory support in patients with COVID-19 (outside intensive care unit). A position paper of the Respiratory Support and Chronic Care Group of the French Society of Respiratory Diseases



C. Rabec^{a,*}, J. Gonzalez-Bermejo^b on behalf the Respiratory Support, Chronic Care Group (GAVO₂) of the French Society of Respiratory Diseases (SPLF)^Δ

^a Service de Pneumologie et Soins Intensifs Respiratoires, Centre Hospitalier Universitaire de Dijon, 14, rue Gaffarel, 21000 Dijon, France

^b Service de Pneumologie, Médecine Intensive et Réanimation (Département R3S, Sorbonne Université, INSERM UMRS 1158 Neurophysiologie respiratoire, expérimentale et clinique, Groupe Hospitalier Universitaire AP-HP Sorbonne Université, AP-HP site Pitié-Salpêtrière, Paris, France

ARTICLE INFO

Article history:

Received 18 April 2020

Received in revised form 9 May 2020

Accepted 10 May 2020

Available online 27 May 2020

ABSTRACT

With first cases noted towards the end of 2019 in China, COVID-19 infection was rapidly become a devastating pandemic. Even if most patients present with a mild to moderate form of the disease, the estimated prevalence of COVID-19-related severe acute respiratory failure (ARF) is 15–20% and 2–12% needed intubation and mechanical ventilation. In addition to mechanical ventilation some other techniques of respiratory support could be used in some forms of COVID-19 related ARF. This position paper of the Respiratory Support and Chronic Care Group of the French Society of Respiratory Diseases is intended to help respiratory clinicians involved in care of COVID-19 pandemic in the rational use of non-invasive techniques such as oxygen therapy, CPAP, non-invasive ventilation and high flow oxygen therapy in managing patients outside intensive care unit (ICU). The aims are: (1) to focus both on the place of each

* Corresponding author.

E-mail address: claudio.rabec@chu-dijon.fr (C. Rabec).

^Δ GAVO₂ collaborators

M. Mercy (Pulmonologist)^c, Metz, France
 L. Grassion (Pulmonologist)^d, Bordeaux, France
 S. Pontier (Pulmonologist)^e, Toulouse, France
 M. Patout (Pulmonologist)^f, Paris, France
 R. Luque (Biomedical Engineer)^g, Jouy-en-Josas, France
 C. Delafosse (Pulmonologist)^h, Eaubonne, France
 C. Raheison-Semjen (Pulmonologist)^d, Bordeaux, France
 B. Maître (Pulmonologist)ⁱ, Créteil, France
 L. Duthoit (Pulmonologist)^j, Lille, France
 A. Mendoza (Respiratory Technician)^f, Paris, France
 L. Jacquin (Respiratory Therapist)^k, Saint-Priest, France
 J.C. Borel (Respiratory Therapist)^l, Grenoble, France
 P. Cervantes (Pulmonologist)^c, Metz, France
 J.-P. Janssens (Pulmonologist)^m, Genève, Switzerland
 J.-F. Chabot (Pulmonologist)ⁿ, Nancy, France
 C. Morelot-Panzini (Pulmonologist)^f, Paris, France
 D. Jaffuel (Pulmonologist)^o, Montpellier, France

<https://doi.org/10.1016/j.resmer.2020.100768>

2590-0412/© 2020 SPLF and Elsevier Masson SAS. All rights reserved.

technique and in describing practical tips (types of devices and circuit assemblies) aimed to limit the risk of caregivers when using those techniques at high risk spreading of viral particles; (2) to propose a step-by-step strategy to manage ARF outside ICU.

The SARS-CoV-2 has been identified as the agent of the pandemic known as Coronavirus disease 2019 (COVID-19). The first cases were noted towards the end of 2019 in Wuhan, China.

COVID-19 infection is spontaneously resolvable in most cases. The clinical presentation can vary from mild respiratory symptoms to severe pneumonia progressing to fulminant acute respiratory failure (ARF) [1]. COVID-2019 pneumonia is characterized by bilateral infiltrates, which can progress to diffuse alveolar condensations. In less severe patients, computed tomography (CT) shows bilateral ground glass sub pleural opacities [2].

The estimated prevalence of COVID-19-related acute respiratory failure (ARF) is 15–20% [1,3,4]. In different published series, 41% had received O₂, 4–13% of patients non-invasive ventilation (NIV) and 2–12% needed intubation and mechanical ventilation [1,3,5–7].

In addition to MV some other techniques of respiratory support such as non-invasive ventilation (NIV), continuous positive airway pressure (CPAP) or high flow oxygen therapy (HFOT) could be used in some forms of ARF. Those techniques are currently applied in ICU but also in a pulmonary department. But the fact that this pandemic could go beyond the capacity of the health system, may lead that those techniques might be applied in less specialized services. These position paper is intended to help respiratory clinicians involved in care of COVID-19 pandemic in managing patients outside ICU. They cannot be regarded as recommendations and reflect only authors experience during COVID-19 pandemics, their expertise in the field of respiratory support and their critical review of the literature.

1. Available tools

1.1. Oxygen therapy

Oxygen should be used in case of severe COVID-19 pneumonia probably as soon as SpO₂ < 92% with a SpO₂ target between 92 and 96%.

There are no randomized controlled studies concerning O₂ in patients with COVID-19 but it is possible to extrapolate data from studies in severe ARF. A meta-analysis of 25 randomized controlled studies has shown that a strategy with no upper limit on O₂ flow (“liberal” approach) increases the risk of in-hospital death compared to a conservative (“targeted” approach) [8]. This contrast with the results of a recent randomized controlled trial (RCT) comparing a “liberal” O₂ therapy strategy (target SpO₂ > 96%) to a conservative one (target SpO₂ > 88% < 92%). This study showed no difference in 28-days survival but an over mortality at 90 days in the conservative group [9]. Thus, based on those data, a target of > 92 < 96% seems reasonable [10]. There are no controlled studies concerning the better interface to deliver O₂. Based on clinical recommendations, a nasal cannula should be used for mild hypoxia and, if an oxygen flow > 6 L min⁻¹ is needed, a switch for a simple face mask set to deliver 5 to 10 L min⁻¹ should be proposed. A non-rebreather mask between 10 and 15 L min⁻¹ should be used in patients remaining hypoxemic despite using a simple mask. These latter interfaces can ensure FiO₂ of 35–55% and 80–95% respectively FiO₂ depending on

flow and breathing pattern [11]. If available, it could be appropriate to use masks with filtered exhalation port (Filtamask™ or similar). Whatever the interface chosen, patient’s comfort and decrease caregiver’s risk will be prioritized when choosing the interface.

1.2. Continuous Positive Airway Pressure (CPAP)

CPAP with added O₂ could be used to improve oxygenation, if conventional O₂ failed and there is no urgent indication for intubation or as a surrogate while waiting for intubation.

This solution is simpler, less expensive and possibly less harmful than NIV.

CPAP must be applied under a strict supervision of a trained physician. ICU team must be prevented and readily available.

The interfaces used to deliver CPAP must be those available or those more familiar for the team. A helmet could be an alternative to limit exposition to droplets dispersion, and remains an option for expert teams. Circuits and masks must be adapted to reduce caregiver’s risks (Fig. 1).

In hypoxemic ARF, applying an extrinsic positive end expiratory pressure (PEEP) increases alveolar recruitment and improves oxygenation. Furthermore, there is high level of evidence about the efficacy of adding PEEP during invasive ventilation in ARDS patients. Nevertheless, we lack information about the effectiveness of applying non-invasive PEEP. Experiences from Italian and Chinese teams are shared but not published.

As CPAP is easier to use than NIV, more available and need less expertise, this technique could be offered as a first line therapy in selected patients, as a gap to invasive MV, in particular when resources are limited or if there is no immediate access to invasive ventilation. A detail of CPAP-delivering devices, their limits and advantages is showed in Fig. 2.

1.3. Non Invasive Ventilation (NIV)

NIV with added O₂ could be used to improve oxygenation and/or providing ventilatory support, if conventional O₂ failed and there is no urgent indication for intubation or as a surrogate while waiting for intubation.

In those cases NIV must be applied under a strict supervision of a trained physician. ICU team must be prevented and readily available.

NIV should be considered in patients who will not be admitted to ICU for intubation.

The interfaces used to deliver NIV must be those available or those more familiar for the team. A helmet could be an alternative to limit exposition to droplets dispersion, and remains an option for expert teams.

Circuits and masks must be adapted to reduce caregiver’s risks (Fig. 2).




Fig. 1. Different ventilator circuit assemblies to reduce viral spreading. (a) Ventilator using double circuit with an integrated expiratory valve. The filter* must be interposed between the expiratory arm and the ventilator. (b) Ventilator using single-limb circuit with an active expiratory valve: the filter* must be interposed between the mask and the expiratory valve. (c) Ventilator using single limb circuit with intentional leaks. Not-vented mask must be preferred if available. In this case a deported exhalation port (Whisper Swivel or similar) must be added. The filter* must be interposed between the mask and the deported exhalation port. If a non-vented mask is not available an alternative is to seal the intentional leak of the vented mask. In the last case caution must be taken not to block anti-asphyxia valve. I: inspiratory arm; E: expiratory arm. * There are not published studies comparing the efficacy of different filters.

<p>ICU ventilator</p>	<p>Life support ventilator, (single ou double limb circuit/expiratory valve)</p>	<p>Bilevel ventilator (single limb circuit with intentional leak)</p>	<p>Boussignac CPAP</p>
<p>Avantages</p> <ul style="list-style-type: none"> - Possibility to ensure FiO₂ 100% - Monitoring capabilities - Less viral spreading - Deliver EPAP even at low O₂ flow - Adjustable EPAP level - Powerful blower able to maintain EPAP level even with important leaks or high inspiratory effort 	<p>Avantages</p> <ul style="list-style-type: none"> - Less viral spreading - Deliver EPAP even at low O₂ flow - Adjustable EPAP level - Powerful blower able to maintain EPAP level even with important leaks or high inspiratory effort - Internal battery (transport possible) 	<p>Avantages</p> <ul style="list-style-type: none"> - Easier to use and easily available - Deliver EPAP even at low O₂ flow - Adjustable EPAP level - Powerful blower able to maintain EPAP level even with important leaks or high inspiratory effort 	<p>Avantages</p> <ul style="list-style-type: none"> - Easier to use - Doesn't need outlet power supply - Doesn't need settings - Disposable material
<p>Inconvenients</p> <ul style="list-style-type: none"> - Limited availability (only in ICU) - Burdensome for a non-trained physician - No internal battery, needs outlet power supply. Non-transportable 	<p>Inconvenients</p> <ul style="list-style-type: none"> - Limited availability - Burdensome for a non-trained physician 	<p>Inconvenients</p> <ul style="list-style-type: none"> - Should not be used with a vented mask with incorporated leak (viral spread) (see figure 2) - Need complex assembly 	<p>Inconvenients</p> <ul style="list-style-type: none"> - High risk spreading of viral particles (high flow) - Needs a flow of at least 30l/mn to provide a CPAP level of 10 cm H₂O - No available for transport - CPAP level not adjustable without manometer

Fig. 2. Different devices providing non-invasive CPAP therapy.

There are no published data concerning NIV use outside ICU in patients with COVID-19. NIV indications in ICU patients with COVID-19-related ARF range from 11 to 34% [12]. Meta-analysis of RCT conducted in severe hypoxemic ARF showed that NIV could reduce the rate of intubation and mortality. However, they included patients with immunosuppression, acute heart failure (AHF) and postoperative ARF [7,13–15]. Moreover, in ARF etiologies other than AHF, NIV has shown a high level of failure with higher mortality (28%) than those treated with O₂ (23%) or HFOT (13%) [16],

In a cohort of patients with Middle East respiratory syndrome, NIV was associated with better survival and a shorter length of stay (LOS) compared patients who were intubated without previous NIV. However, NIV showed a high failure rate (92.4%) requiring intubation [17]. Finally, associating NIV and prone position reduced intubation in patients with moderate ARDS related to viral pneumonia [18].

On the other side, NIV could be harmful in those patients as it could worsen lung damage due to high pressures and high VT and

could delay intubation [19,20]. Moreover, NIV as CPAP is at high risk spreading of viral particles. In this field, WHO guidelines for the management of ARF in COVID-19 advocate the use of CPAP or NIV, provided that appropriate personal protective equipment is worn [21].

Hence NIV could be considered as a first line therapy outside the ICU in particular when resources are limited or if there is no immediate access to invasive ventilation. In those cases NIV must be applied under a strict supervision of a trained physician to detect early and rapid worsening, a frequent condition in COVID-19 pneumonia. ICU team must be prevented and readily available. When a patient is treated with NIV, monitoring should be intensified. If hypoxemia worsens the patient must be transferred to ICU if eligible for intubation. This decision will integrate not only the clinical severity, but also underlying pathologies and the “living wills” or end-of-life decisions (if available) of the patient and his family. This requires previous discussion on a case-by-case basis.

NIV should also be considered in patients no eligible to be admitted to ICU or in those with do not intubate (DNI) decision. In those cases, NIV must be continued only if it is well tolerated and a benefit is obtained. Otherwise, comfort measures only, including pharmacological measures, may be applied to relieve dyspnea [22].

Some patients with underlying respiratory disease (COPD, obesity hypoventilation, restrictive disease) complicated with COVID-19 pneumonia could benefit from de novo NIV.

In patients on long term NIV, treatment must be adapted during hospitalization, possibly by temporarily lowering the pressures if excessive leaks are present. Circuits and masks must be adapted (Fig. 2).

When available a helmet interface could be applied. A randomized controlled study conducted in ARDS has shown that NIV when delivered by a helmet decrease intubation rate and mortality [23]. Moreover helmet limits viral spreading [24].

It is imperative not to insist with NIV or O₂ in patients that worsen. This can lead to a delay in intubation, which can be fatal. Increased vigilance is necessary since muscle exhaustion is expressed late in those patients. In all cases, the intubation must be anticipated, carried out according to rigorous procedures and under conditions limiting the risk of caregivers.

As NIV is at high risk spreading of viral particles circuits and masks must be adapted to reduce caregiver's risks [14,25,26] (Fig. 2). Moreover, caution may be taken to ensure a good interface fitting for CPAP or NIV systems, to minimize widespread dispersion of exhaled air and reduce risk of airborne transmission.

1.4. High Flow Oxygen Therapy (HFOT)

HFOT could be used to improve oxygenation, if conventional O₂ failed and there is no urgent indication for intubation or as a surrogate while waiting for intubation.

As HFOT is at high risk spreading of viral particles, protective measures must be applied to reduce caregiver's exposition.

HFOT delivers a high-flow gas mixture (up to 70L/min) with variable FiO₂ (up to 100%) administered by a nasal cannula. Compared to conventional oxygen HFOT can ensure a constant and known FiO₂. Other advantages are dead space reduction and generation of a low PEEP level allowing alveolar recruitment [27].

There are no published data concerning HFOT in patients with COVID-19 pneumonia.

In a RCT conducted in severe hypoxemic ARF, HFOT reduced 90-days mortality but not intubation rate compared to conventional oxygen [17]. Otherwise, a meta-analysis of 9 RCT shows a decrease in intubation rate but without improving survival or length of stay [28–30] an important issue in the field of COVID-19 pandemic considering an expected scarcity of ventilators.

As with NIV, adding prone position to HFOT may help to avoid intubation in patients with moderate ARDS related to viral pneumonia [18].

The same precautions cited above for NIV should be taken if HFOT is used as a first line therapy in patients qualifying for intubation.

Even if published studies did not demonstrate an increased risk for caregivers when comparing HFOT to conventional O₂ [31,32] to limit the risk of contamination, the following measures are suggested when using HFOT:

- ensure maximum sealing of the interface;
- limit the flow rate to the minimum necessary. Prefer higher FiO₂ instead of higher flow (i.e.: start at 30L/min and increase FiO₂ to ensure target SaO₂);
- patient must wear a surgical mask during care.

A detail of HFOT-delivering devices, their limits and advantages is showed in Fig. 3.

1.5. Nebulized treatments

As nebulization is at high risk spreading of viral particles, nebulized treatments should be limited as much as possible.

Spray and powders should be preferred to provide inhaled therapy, preferably by using a personal inhalation chamber. If not possible, minimal distance of 1 m must be respected and the room must aerate during nebulization.

1.6. Tracheal aspirations

In the case of tracheostomized ventilator-dependent patients a “closed suction system” must be used.




Fig. 3. Different devices providing HFOT therapy.

2. Patient management

The proposed flow chart strategy to manage ARF outside ICU in COVID-19 patients is showed in Fig. 4.

It is crucial to ensure an appropriate triage of patients at the admission. As those patients are at risk of early and rapid worsening, it seems reasonable that those with a rapidly progressive form (i.e presenting a SpO₂ < 94% and a RR > 30 while receiving > 6 L O₂) must be promptly assessed by the ICU staff to decide proper allocation.

2.1. Step by step approach

- Check and treat comorbidities.
- Begin conventional oxygen therapy by using nasal prongs to obtain the target RR and SpO₂:
 - if predominant oral breathing or intolerance to high flow, an oxygen mask could be proposed,
 - masks with filtered exhalation port could be used if available,
 - there is no limit in terms of O₂ flow rate but if > 6 L/min or a mask with reservoir bag is needed, ICU team must be prevented and promptly available,
 - venturi masks should be precluded;
- then, if oxygen therapy failed, non-invasive respiratory assistance should be proposed with CPAP and/or HFOT as a first choice. NIV should be used as a second line therapy in case of CPAP or HFOT failure, and mainly if hypercapnia develops;
- HFOT could be an alternative in the absence of CPAP/NIV or as a therapeutic ceiling option (HFOT allows higher FiO₂ but there is hypothetically a greater risk of drops diffusion and low PEEP levels are generated) [33].

- Close monitoring is needed during at least the first 48–72 hours, including SpO₂, RR and clinical assessment (ventilatory mechanics/use of accessory muscles). Three issues highlighted by different teams managing those patients requires particular attention:
 - an initially stable patient may suddenly become unstable (with refractory hypoxemia and high fever),
 - a delayed re-aggravation was noted in a significant percentage of patients (stability then rapid worsening after 48 h, up to 7 days),
 - it was described in some of these patients an impaired perception of dyspnea despite severe hypoxemia. A possible explanation is a neuroinvasive potential of COVID-19 that could spread in brainstem affecting respiratory center and/or mechano-/chemoreceptors [34].
- In cases with DNI decision, it is recommended to dispose an end-of-life sedation protocol to be applied if patient's condition worsens (<http://www.sfap.org/actualite/outils-et-ressources-soins-palliatifs-et-covid-19>).

In summary: non- invasive respiratory support could be useful in treating COVID-19-related ARF. A rational use of different techniques (oxygen therapy, CPAP, NIV or HFOT) by a trained pulmonologist could allow to prevent clinical aggravation and reduce the risk of ICU admission. Then, those techniques should be considered as a first line therapy outside the ICU in particular when resources are limited or if there is no immediate access to invasive ventilation. A step-by step approach, under a strict supervision of a trained physician, must be applied to detect early and rapid aggravation. In patients eligible for intubation. ICU team must be prevented and readily available to an immediate transfer to ICU as soon as the patient's condition impairs.




Fig. 4. Flow chart strategy for managing ARF in COVID-19 patients (modified from [33]) ABG: Arterial Blood Gases; COPD: chronic obstructive pulmonary disease; RR: respiratory rate; HFOT: high flow oxygen therapy; CPAP: continuous positive airway pressure; NIV: non-invasive ventilation; EPAP: expiratory positive airway pressure; EI: endotracheal intubation).

Disclosure of interest

The authors have not supplied their declaration of competing interest.

Acknowledgements

Stefano Nava (Italy), Javiers Sayas (Spain), Michelle Chatwin (UK), Manel Lujan (Spain), Annalisa Carlucci (Italy) for sharing information.

Special acknowledgement to the colleagues of the Italian Thoracic Society (AIPO-ITS) and Italian Respiratory Society (SIP/IRS) for their original idea of the flow chart proposed in this paper.

A French version of this document is available online in open format (<http://www.splf.fr/wp-content/uploads/2020/04/RespiPreREA-SPLF-GAVO2avrii2020.pdf>).

References

- [1] Wu Z, McGoogan JM. Characteristics of and important lessons from the Coronavirus Disease 2019 (COVID-19) outbreak in China: summary of a report of 72,314 cases from the Chinese center for disease control and prevention. *JAMA* 2020.
- [2] Wang Y, Dong C, Hu Y, Li C, Ren Q, Zhang X, et al. Temporal changes of CT findings in 90 patients with COVID-19 pneumonia: a longitudinal study. *Radiology* 2020:200843.
- [3] Guan W-J, Ni Z-Y, Hu Y, Liang W-H, Ou C-Q, He J-X, et al. Clinical characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020.
- [4] Shi Y, Yu X, Zhao H, Wang H, Zhao R, Sheng J. Host susceptibility to severe COVID-19 and establishment of a host risk score: findings of 487 cases outside Wuhan. *Crit Care* 2020;24:108.
- [5] Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;395:507–13.
- [6] Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel Coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020.
- [7] Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 2020.
- [8] Chu DK, Kim LH-Y, Young PJ, Zamiri N, Almenawer SA, Jaeschke R, et al. Mortality and morbidity in acutely ill adults treated with liberal versus conservative oxygen therapy (IOTA): a systematic review and meta-analysis. *Lancet* 2018;391:1693–705.
- [9] Barrot L, Asfar P, Mauny F, Winiszewski H, Montini F, Badie J, et al. Liberal or conservative oxygen therapy for acute respiratory distress syndrome. *N Engl J Med* 2020;382:999–1008.
- [10] Alhazzani W, Møller MH, Arabi YM, Loeb M, Gong MN, Fan E, et al. Surviving sepsis campaign: guidelines on the management of critically ill adults with Coronavirus Disease 2019 (COVID-19). *Crit Care Med* 2020, <http://dx.doi.org/10.1097/CCM.0000000000004363> [Epub ahead of print].
- [11] O'Driscoll B, Howard LS, Earis J, Mak V. BTS guideline for oxygen use in adults in healthcare and emergency settings. *Thorax* 2017;72:i1–90.
- [12] Grasselli G, Pesenti A, Ceconi M. Critical care utilization for the COVID-19 outbreak in Lombardy, Italy: early experience and forecast during an emergency response. *JAMA* 2020 [Epub 2020/03/14].
- [13] Wang T, Zhang L, Luo K, He J, Ma Y, Li Z, et al. Noninvasive versus invasive mechanical ventilation for immunocompromised patients with acute respiratory failure: a systematic review and meta-analysis. *BMC Pulmon Med* 2016;16:129.
- [14] Xu X-P, Zhang X-C, Hu S-L, Xu J-Y, Xie J-F, Liu S-Q, et al. Noninvasive ventilation in acute hypoxemic nonhypercapnic respiratory failure: a systematic review and meta-analysis. *Crit Care Med* 2017;45:e727–33.
- [15] Zayed Y, Banifadel M, Barbarawi M, Kheiri B, Chahine A, Rashdan L, et al. Non-invasive oxygenation strategies in immunocompromised patients with acute hypoxemic respiratory failure: a pairwise and network meta-analysis of randomized controlled trials. *J Intensive Care Med* 2019 [885066619844713].
- [16] Frat J-P, Thille AW, Mercat A, Girault C, Ragot S, Perbet S, et al. High-flow oxygen through nasal cannula in acute hypoxemic respiratory failure. *N Engl J Med* 2015;372:2185–96.
- [17] Alraddadi BM, Qushmaq I, Al-Hameed FM, Mandourah Y, Almekhlafi GA, Jose J, et al. Noninvasive ventilation in critically ill patients with the Middle East respiratory syndrome. *Influenza Other Respir Viruses* 2019;13:382–90.
- [18] Ding L, Wang L, Ma W, He H. Efficacy and safety of early prone positioning combined with HFNC or NIV in moderate to severe ARDS: a multi-center prospective cohort study. *Crit Care* 2020;24:28.
- [19] Brochard L, Lefebvre J-C, Cordioli RL, Akoumianaki E, Richard J-CM. Noninvasive ventilation for patients with hypoxemic acute respiratory failure. *Semin Respir Crit Care Med* 2014;35:492–500.
- [20] Slutsky AS, Ranieri VM. Ventilator-induced lung injury. *N Engl J Med* 2013;369:2126–36.
- [21] WHO Clinical management of severe acute respiratory infection when novel coronavirus (2019-nCoV) infection is suspected: interim guidance; 2020 <https://www.apps.who.int/iris/handle/10665/330893>.
- [22] Tripodoro VA, Rabec CA, De Vito EL. Withdrawing noninvasive ventilation at end-of-life care: is there a right time? *Curr Opin Support Palliat Care* 2019;13:344–50.
- [23] Patel BK, Wolfe KS, Pohlman AS, Hall JB, Kress JP. Effect of noninvasive ventilation delivered by helmet vs face mask on the rate of endotracheal intubation in patients with acute respiratory distress syndrome: a randomized clinical trial. *JAMA* 2016;315:2435–41.
- [24] Hui DS, Chow BK, Lo T, Ng SS, Ko FW, Gin T, et al. Exhaled air dispersion during noninvasive ventilation via helmets and a total facemask. *Chest* 2015;147:1336–43.
- [25] Fowler RA, Guest CB, Lapinsky SE, Sibbald WJ, Louie M, Tang P, et al. Transmission of severe acute respiratory syndrome during intubation and mechanical ventilation. *Am J Respir Crit Care Med* 2004;169:1192–202.
- [26] Hui DSC, Zumla A. Severe acute respiratory syndrome: historical, epidemiologic, and clinical features. *Infect Dis Clin N Am* 2019;33:869–89.
- [27] Drake MG. High-flow nasal cannula oxygen in adults: an evidence-based assessment. *Ann Am Thorac Soc* 2018;15:145–55.
- [28] Ni Y-N, Luo J, Yu H, Liu D, Liang B-M, Liang Z-A. The effect of high-flow nasal cannula in reducing the mortality and the rate of endotracheal intubation when used before mechanical ventilation compared with conventional oxygen therapy and noninvasive positive pressure ventilation. A systematic review and meta-analysis. *Am J Emerg Med* 2018;36:226–33.
- [29] Ou X, Hua Y, Liu J, Gong C, Zhao W. Effect of high-flow nasal cannula oxygen therapy in adults with acute hypoxemic respiratory failure: a meta-analysis of randomized controlled trials. *CMAJ* 2017;189:E260–7.
- [30] Rochweg B, Granton D, Wang DX, Helviz Y, Einav S, Frat JP, et al. High flow nasal cannula compared with conventional oxygen therapy for acute hypoxemic respiratory failure: a systematic review and meta-analysis. *Intensive Care Med* 2019;45:563–72.
- [31] Leung CCH, Joynt GM, Gomersall CD, Wong WT, Lee A, Ling L, et al. Comparison of high-flow nasal cannula versus oxygen face mask for environmental bacterial contamination in critically ill pneumonia patients: a randomized controlled crossover trial. *J Hosp Infect* 2019;101:84–7.
- [32] Raboud J, Shigayeva A, McGeer A, Bontovics E, Chapman M, Gravel D, et al. Risk factors for SARS transmission from patients requiring intubation: a multicentre investigation in Toronto, Canada. *PLoS One* 2010;5:e10717. <http://www.aiponet.it/news/speciale-covid-19/2426-managing-the-respiratory-care-of-patients-with-covid-19-english-version.html> [updated 2020-04-01 21:35:40].
- [34] Li Y-C, Bai W-Z, Hashikawa T. The neuroinvasive potential of SARS-CoV2 may play a role in the respiratory failure of COVID-19 patients. *J Med Virol* 2020.